

Thermodynamics of Large- N Super Yang-Mills Theory and AdS/CFT Correspondence ¹

Chanju Kim^a and Soo-Jong Rey^{b,c}

^a *School of Physics, Korea Institute for Advanced Study, Seoul 702-701 Korea*

^b *Physics Department, Seoul National University, Seoul 151-742 Korea*

^c *Asia-Pacific Center for Theoretical Physics, Seoul 130-012 Korea*

cjkim@ns.kias.re.kr, sjrey@gravity.snu.ac.kr

abstract

Thermodynamics of $d = 4$, $\mathcal{N} = 4$ supersymmetric $SU(N)$ Yang-Mills theory is studied with particular attention on perturbative expansion at weak 't Hooft coupling regime and interpolation to strong coupling regime thereof. Non-ideal gas effect to free-energy is calculated and found that leading- and next-to-leading-order corrections contribute with relative opposite signs. Padé approximant method is adopted to improve fixed-order, perturbative series and is found to decrease free-energy monotonically as 't Hooft coupling parameter is increased. This may be regarded as an indication of smooth interpolation of thermodynamics between weak and strong 't Hooft coupling regimes, as suggested by Maldacena's AdS/CFT correspondence.

¹Work supported in part by KOSEF Interdisciplinary Research Grant 98-0702-07-01-5 and SRC Program, KRF International Collaboration Grant, Ministry of Education Grant 98-015-D00054, and The Korea Foundation for Advanced Studies Faculty Fellowship.

1 Introduction

Maldacena's proposal [1] relates large N limit of the $d = 4$, $\mathcal{N} = 4$ supersymmetric $SU(N)$ Yang-Mills theory to Type IIB superstring theory on $AdS_5 \times S^5$. According to the correspondence, parameters of Yang-Mills and string theories are related each other as:

$$g_{\text{YM}}^2 = 2\pi g_{\text{st}}, \quad g_{\text{YM}}^2 N = \frac{1}{2} \frac{R^4}{\alpha'^2},$$

where R is the curvature radii of both AdS_5 and S^5 , and g_{st} and α' are string coupling parameter and string slope parameter, respectively. Exploration of the proposed duality over the entire parameter space (g_{YM}^2, N) is seemingly a formidable, if not impossible, task. However, the above relation already implies that physics of super Yang-Mills theory in the strong 't Hooft coupling limit, $\lambda^2 \equiv g_{\text{YM}}^2 N \rightarrow \infty$, is directly accessible from systematic expansion in the zero-slope limit, $(\alpha'/R^2) \rightarrow 0$, of Type IIB string theory, viz. gauged AdS_5 supergravity. Indeed, all the checkpoints regarding the duality examined so far have been exclusively in this limit. It is therefore of interest to explore, for various physical quantities, the Maldacena's proposal beyond the leading order in strong 't Hooft coupling in super Yang-Mills theory and in (α'/R^2) in gauged AdS_5 supergravity. Of particular interest is whether the physical quantity is an analytic function over the entire parameter space. If so, it would be interpolating smoothly between weak and strong coupling regimes. In this paper, as an example of such physical quantities, we would like to investigate thermodynamic free-energy density $F(N, \lambda)$, with particular attention to the issue of interpolation between weak and strong 't Hooft coupling regimes. The issue is particularly relevant, as, based on certain assumptions on analyticity and convergence of the perturbative series, it has been claimed that such a smooth interpolation is not possible [2].

In the strong 't Hooft coupling limit, the free-energy density has been calculated in [3], including leading-order correction. Their calculation goes as follows. According to the Maldacena's proposal extended to finite temperature, super Yang-Mills theory at high temperature T is described, in large N and strong 't Hooft coupling limit, by a Schwarzschild AdS_5 black hole [4] at Hawking temperature T . The metric of the black hole is given by

$$ds^2 = \frac{U^2}{R^2} \left(-f(U) dt^2 + d\mathbf{x}_{\parallel}^2 \right) + \frac{R^2}{U^2} \left(f^{-1}(U) dr^2 + U^2 d\Omega_5^2 \right), \quad (1)$$

where, expressed in terms of Yang-Mills parameters, the Schwarzschild factor $f(U)$ is given by

$$f(U) = \left(1 - \frac{U_0^4}{U^4} \right), \quad U_0 = 2\pi\lambda T.$$

As shown in [3], the metric Eq.(1) turns out to be a solution of gauged AdS_5 supergravity equations of motion not only at leading-order (Einstein-Hilbert action) but also including $\mathcal{O}(\alpha'^3)$

string correction terms. The free-energy density F is then obtained from the Euclidean supergravity action of the black hole times Hawking temperature T . The result is

$$\begin{aligned} F(N, T) &= h(\lambda) \left(-\frac{1}{6} \pi^2 N^2 T^4 \right) \\ h(\lambda) &= \left[\frac{3}{4} + \frac{45}{32} \zeta(3) \lambda^{-3/2} + \dots \right], \end{aligned} \quad (2)$$

and hence indicates that the free-energy density is diminished by a factor $h(\lambda) \sim 3/4$ compared to the free-energy of ideal gas of $\mathcal{N} = 4$ super Yang-Mills theory, $-\pi^2/6 \cdot N^2 T^4$. Branch-cut of the free-energy density at $\lambda = \infty$ is a genuine prediction for the strong coupling limit, along with the heavy quark potential, both at zero [5, 6] and finite temperatures [7, 8, 9], and anomalous dimensions of long supermultiplets [10, 11].

Were it interpolating smoothly over the entire 't Hooft coupling parameter space, the function $h(\lambda)$ in Eq.(2) is expected to make monotonic cross-over between $h(\lambda \rightarrow 0) \sim 1$ and $h(\lambda \rightarrow \infty) \sim 3/4$. To examine whether this is the case, in this paper, we calculate the free-energy density of the super Yang-Mills theory in fixed-order perturbative expansion and examine its behavior as λ is increased. According to our calculation, the leading-order $\mathcal{O}(\lambda^2)$ correction indeed makes $h(\lambda)$ decreasing, but the next-order, non-analytic $\mathcal{O}(\lambda^3)$ correction contributes with (relative) opposite sign. The result up to this order then seems not to exhibit anticipated monotonically interpolating behavior, but it may have to do with truncation of the perturbation expansion to lower order terms. In an attempt to improve the fixed-order perturbation theory and (partially) resum higher-order terms, we have employed Padé approximant method. Our result shows that, once Padé-improved, the function $h(\lambda)$ indeed becomes monotonically decreasing as it is extrapolated to larger values of the 't Hooft coupling parameter. Interestingly enough, the (relative) opposite sign of the $\mathcal{O}(\lambda^3)$ correction turns out precisely the origin of the monotonically decreasing behavior after the Padé approximant is made.

The paper is organized as follows. In section 2, we present perturbative calculation of the free-energy density of $d = 4$, $\mathcal{N} = 4$ super Yang-Mills theory up to $\mathcal{O}(\lambda^3)$. In section 3, we improve the fixed-order perturbative result of section 2 by Padé approximant method and find smooth interpolation plausible. In section 4, we conclude with brief discussion on interpolation of free-energy density between $d = 2$, $\mathcal{N} = 4$ supersymmetric $SU(N_1) \times SU(N_5)$ Yang-Mills theory and Type IIB supersring theory on AdS_3 . Preliminary results of the present paper has been announced by one of the authors (SJR) in [12]. Since then, while our paper is being prepared, two independent works dealing with aspects of the calculation in section 2 up to $\mathcal{O}(\lambda^2)$ [13] and $\mathcal{O}(\lambda^3)$ [14] have been reported. Wherever overlapping, our results are in complete agreement with theirs.

2 Perturbative Calculation of Free-Energy

In this section, we will be computing free-energy of four-dimensional super Yang-Mills theory with $\mathcal{N} = 4$ supersymmetry and gauge group $G = SU(N)$ up to $\mathcal{O}(g_{\text{YM}}^3)$ in large- N limit. Throughout the computation, we will adopt Feynman gauge and use dimensional regularization for ultraviolet divergences. In component fields, the $\mathcal{N} = 4$ super Yang-Mills theory consists of a gauge boson, four Majorana fermions and three complex scalars in the adjoint representation of the gauge group G .




Figure 1: *One-loop Feynman diagrams contributing to free-energy F_1 . Gauge bosons are represented by wiggly lines, ghosts by dotted lines, fermions by solid lines, and scalars by dashed lines.*

At leading order, $\mathcal{O}(g_{\text{YM}}^0)$, the free energy density F_1 is simply that of the non-interacting massless degrees of freedom and is obtained from one-loop Feynman diagrams, as shown in Fig.1 for gauge bosons, ghosts, gaugino and three adjoint fermions, and three adjoint scalars, respectively. Evaluating each diagram explicitly, one obtains

$$-F_1(G, T) = -4d_G \cdot \frac{1}{2}\mathcal{B}_0 + 2d_G \cdot \frac{1}{2}\mathcal{B}_0 + 4d_G \cdot 2 \cdot \frac{1}{2}\mathcal{F}_0 - 3d_G \cdot 2 \cdot \frac{1}{2}\mathcal{B}_0.$$

Here, d_G denotes the dimension of the gauge group G , and $\mathcal{B}_0(T)$ and $\mathcal{F}_0(T)$ are defined as²

$$\begin{aligned}\mathcal{B}_0(T) &= \sum_{[p]} \ln p^2 = -\frac{\pi^2}{45}T^4, \\ \mathcal{F}_0(T) &= \sum_{\{p\}} \ln p^2 = \frac{7}{8} \cdot \frac{\pi^2}{45}T^4.\end{aligned}$$

²Here, $[p]$ ($\{p\}$) represents that the four-momentum $p = (p_0, \mathbf{p})$ is bosonic (fermionic), namely,

$$\begin{aligned}\sum_{[p]} &\equiv T \sum_{p_0} \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \quad \text{where} \quad p_0 = 2\pi n T, \\ \sum_{\{p\}} &\equiv T \sum_{p_0} \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \quad \text{where} \quad p_0 = 2\pi(n + \frac{1}{2})T.\end{aligned}$$

Hence, summing up all the one-loop diagrams, one obtains $F_1(G, T)$ as

$$\begin{aligned} F_1(G, T) &= 4d_G (\mathcal{B}_0 - \mathcal{F}_0) \\ &= -\frac{\pi^2}{6} d_G T^4. \end{aligned} \quad (3)$$

The contribution of the next-leading $\mathcal{O}(g_{\text{YM}}^2)$ order is calculated from two-loop Feynman diagrams in Fig. 2.




Figure 2: Two-loop diagrams contributing to the free-energy density F_2 .

After a straightforward calculation, one finds that each diagram gives rise to the following contribution:

$$\begin{aligned} -F_{2a} &= -3g_{\text{YM}}^2 c_A d_G \mathcal{B}^2 \\ -F_{2b} &= +\frac{9}{4} d_G g_{\text{YM}}^2 c_A d_G \mathcal{B}^2 \\ -F_{2c} &= -\frac{1}{4} g_{\text{YM}}^2 c_A d_G \mathcal{B}^2 \\ -F_{2d} &= -16g_{\text{YM}}^2 c_A d_G (\mathcal{B}^2 - 2\mathcal{B}\mathcal{F}) \\ -F_{2e} &= -\frac{15}{2} g_{\text{YM}}^2 c_A d_G \mathcal{B}^2 \\ -F_{2f} &= -12g_{\text{YM}}^2 c_A d_G \mathcal{B}^2 \\ -F_{2g} &= +\frac{9}{2} g_{\text{YM}}^2 c_A d_G \mathcal{B}^2. \end{aligned} \quad (4)$$

Here, $c_A(G)$ denotes the quadratic Casimir invariant of the gauge group G and

$$\mathcal{B}(T) \equiv \sum_{[p]} \frac{1}{p^2} = +\frac{T^2}{12},$$

$$\mathcal{F}(T) \equiv \sum_{\{p\}} \frac{1}{p^2} = -\frac{T^2}{24}.$$

Adding up all the two-loop Feynman diagrams, one finds

$$\begin{aligned} F_2(G, T) &= 16g^2 c_A d_G (\mathcal{B} - \mathcal{F})^2 \\ &= \frac{1}{4} d_G (g_{\text{YM}}^2 c_A) T^4. \end{aligned} \quad (5)$$

Comparison with Eq.(3) shows that the perturbative expansion is indeed in powers of the 't Hooft coupling parameter $\lambda^2 \equiv g_{\text{YM}}^2 c_A$. Note that, at zero temperature, $\mathcal{B}_0 = \mathcal{F}_0$ and $\mathcal{B} = \mathcal{F}$ and hence bosonic and fermionic contributions in Eqs.(3), (5) are cancelled out, manifesting the supersymmetry nonrenormalization theorem for the vacuum energy. For $G = SU(N)$, where $d_G = N^2 - 1$ and $c_A = N$, the results Eqs.(3), (5) are in agreement with [13, 14].

In the fixed-order perturbative expansion, the next order would be of $\mathcal{O}(g_{\text{YM}}^4)$, coming from three-loop Feynman diagrams. However, as is well-known, (a subset of) these diagrams are afflicted with severe infrared divergences associated with the exchange of electrostatic gauge bosons and scalars. Physically, these infrared divergences are screened by Debye plasma screening. The screening of gauge bosons and scalars can be taken into account by a resummation of so-called ring diagrams to all orders [15, 16]. The infinite-order resummation is the origin of the non-analytic contribution starting at order $(g_{\text{YM}}^2)^{3/2}$. An efficient way to calculate the non-analytic term is to reorganize the fixed-order, perturbative expansion by rewriting the Lagrangian as [17]

$$\mathcal{L}_{SYM} = \left(\mathcal{L}_{SYM} + \frac{1}{2} M_{\text{el}}^2 A_0^2 + m_{\text{el}}^2 \sum_{i=1}^3 \phi_i^\dagger \phi_i \right) - \left(\frac{1}{2} M_{\text{el}}^2 A_0^2 + \frac{1}{2} m_{\text{el}}^2 \sum_{i=1}^3 \phi_i^\dagger \phi_i \right),$$

and to treat the second term as perturbative interactions. Here, M_{el} is the electric thermal mass for the gauge boson given by the one-loop self-energy $\Pi_{\mu\nu}$ at zero momentum³, $M_{\text{el}}^2 \delta^{ab} = \Pi_{00}^{ab}(0)$, and m_{el} is the thermal mass for scalars calculated from $m_{\text{el}}^2 \delta^{ab} = \Pi^{ab}(0)$. Self-energy diagrams in Fig. 3 can be evaluated straightforwardly in the zero-momentum limit, much as was done in computation of Eq. (4). The result is

$$\begin{aligned} M_{\text{el}}^2 &= 2 (g_{\text{YM}}^2 c_A) T^2, \\ m_{\text{el}}^2 &= (g_{\text{YM}}^2 c_A) T^2. \end{aligned} \quad (6)$$

Rather than presenting details of straightforward but tedious calculation, , we briefly discuss how $\mathcal{O}(g_{\text{YM}}^3)$ contribution can be obtained. The free-energy density of an ideal Bose gas with

³A nonzero thermal mass is generated for A_0 , but not for A_i . This stems from the fact that, after dimensional reduction over Euclidean time direction, one still retains $d = 3$ gauge invariance.




Figure 3: *One-loop self-energy diagrams for gauge bosons and scalars.*

mass m is given by

$$\begin{aligned}
F_{\text{free boson}} &= \frac{1}{2} \sum_{[p]} \ln(p^2 + m^2), \\
&= -\frac{\pi^2}{90} T^4 + \frac{1}{24} m^2 T^2 - \frac{1}{12\pi} m^3 T + \mathcal{O}(m^4).
\end{aligned} \tag{7}$$

The first term is the well-known, black-body radiation effect of the massless scalar particles. It exerts a positive radiation pressure and hence a negative contribution to the free-energy. The second term is the effect of nonzero mass; it contributes positively to the free energy, as expected on physical ground. The third term, which gives rise to a negative correction, may be understood as representing the bosonic nature of the particle⁴. Its contribution is of $\mathcal{O}(m^3)$, thus cannot be obtained in a naive perturbative expansion in powers of m^2 . If one interprets the mass m as the thermal mass due to the Debye plasma screening, Eq.(6), then Eq.(7) may be understood as an expansion with respect to the coupling $(g_{\text{YM}}^2 c_A)^{1/2}$. In particular, one finds that the sign of the g_{YM}^3 term is negative. Actually, the $\mathcal{O}(g_{\text{YM}}^3)$ term of the free energy comes entirely from Eq.(7) and there is no further correction of order g_{YM}^3 . The reason is

⁴For a Fermi gas, there is no infrared divergence and hence no non-analytic contribution to the free-energy either.

simply because the thermal mass m_{el} is defined in such a way that it precisely reproduces the resummed g_{YM}^3 effect in the form of Eq.(7). Even though there are other g_{YM}^3 order contributions from individual two-loop diagrams Fig. 2, they are all cancelled out by the corresponding mass counterterm one-loop diagrams shown in Fig. 4.



Figure 4: *Mass-counterterm diagrams due to thermal Debye masses.*

Therefore, on general ground, the $\mathcal{O}(g_{\text{YM}}^3)$ contribution due to Debye plasma screening contributes negatively to the free-energy (viz. positive contribution to the radiation pressure), independent of the details of the theory under consideration and can be read from Eq.(7) once the thermal masses are calculated. Incidentally, extra terms of order g_{YM}^2 in Eq.(7) are also cancelled out by the mass counterterms given in Fig. 4, and have no net contribution, as should be so for consistency of the perturbative expansion.

From the above discussion, it is now quite a simple matter to evaluate the free-energy density F_3 of $\mathcal{O}(g_{\text{YM}}^3)$:

$$\begin{aligned} F_3 &= d_G \left(-\frac{1}{12\pi} M_{\text{el}}^3 T \right) - 6d_G \left(-\frac{1}{12\pi} m_{\text{el}}^3 T \right), \\ &= -\frac{3+\sqrt{2}}{12\pi} d_G (g^2 c_A)^{3/2} T^4, \end{aligned} \quad (8)$$

where the relevant degeneracy factors are also taken into account.⁵ Adding Eqs.(3), (5) and (8) altogether, one finally arrives at the free energy density of four-dimensional super Yang-Mills theory with $\mathcal{N} = 4$ and gauge group G up to $\mathcal{O}(g_{\text{YM}}^3)$,

$$F = -\frac{\pi^2}{6} d_G T^4 \left\{ 1 - \frac{3}{2\pi^2} (g_{\text{YM}}^2 c_A) + \frac{3+\sqrt{2}}{\pi^3} (g_{\text{YM}}^2 c_A)^{3/2} + \mathcal{O}((g_{\text{YM}}^2 c_A)^2) \right\}. \quad (9)$$

⁵ One can verify by an explicit calculation that the other terms of order g_{YM}^2 and g_{YM}^3 are all cancelled out.

3 Padé Approximant of Free Energy

The next order correction to Eq.(9) will be of order g_{YM}^4 and can be calculated by considering three-loop diagrams. Moreover, a full account of the Debye screening at three loop order would produce terms of order $g_{\text{YM}}^2 \ln g_{\text{YM}}^2$ and g_{YM}^5 . These contributions can be evaluated by utilizing methods developed in Refs. [17], [18]. On the other hand, extending perturbative evaluation of the free-energy density to even higher orders breaks down beginning at order g_{YM}^6 . This is mainly due to another source of infrared problems associated with exchange of *magnetostatic* gluons. Screening of the magnetostatic gluons cannot be achieved by resummation of the perturbative diagrams and, as the effect is inherently nonperturbative, can only be treated using a reliable method such as lattice simulation [16]. As the new infrared divergence appears beginning at order g_{YM}^6 , by scaling analysis, the *magnetic* screening mass scale is of order $g_{\text{YM}}^2 T$.

Rather than dwelling on estimates of these effects further, one might like to explore implication of the result Eq.(9) to the behavior of the perturbative free-energy as the 't Hooft coupling parameter λ becomes large. In particular, one would like to study if the free-energy at weak coupling regime Eq.(9) may possibly be interpolated monotonically to that at strong coupling regime Eq.(2) obtained via AdS_5 supergravity. On a first look, as λ is increased, the positive sign of g_{YM}^3 term seems to indicate that the result Eq.(9) at weak coupling regime behaves against any smooth interpolation to strong coupling regime. However, it might well have to do with the nature of the fixed-order perturbation theory, which truncates physical quantities to finite-order polynomials and induce fluctuating behavior. Certainly, if possible at all, one needs to improve the perturbative result Eq.(9) before any definite conclusion is drawn.

Replacement of the fixed-order perturbation series by Padé approximant [19] may be considered as a simple tool of gaining an idea on how the series would behave as the coupling parameter is increased. Indeed, the Padé approximant of QCD and other field theories, which has been applied already to high-energy processes [20] quite successfully ⁶, seems to improve convergence of the perturbative free-energy in the right direction [21]. We will thus apply the Padé's approximant to the perturbative free-energy density of the $\mathcal{N} = 4$ super Yang-Mills theory and examine whether the approach lead to monotonic interpolation between weak and strong coupling regimes.

Let us write Eq. (9) as

$$R(\lambda) \equiv \frac{F(\lambda, T)}{F(\lambda = 0, T)} = 1 + f_2 \lambda^2 + f_3 \lambda^3 + \cdots, \quad (10)$$

where, from Eq.(9), $f_2 = -3/2\pi^2$ and $f_3 = (3 + \sqrt{2})/\pi^3$, respectively. Note that, as it stands, $R(\lambda)$ exhibits a minimum at $\lambda = (-2f_2/3f_3)$, viz. $g_{\text{YM}}^2 N \approx 0.5$. The general form of the Padé

⁶in the sense that Padé approximant "postdicts" the known higher-order perturbative terms and also reduces renormalization scale and scheme dependences.

approximants to Eq.(10) is given by

$$R_{[N,L]}(\lambda) = \frac{1 + \sum_{n=1}^N c_n \lambda^n}{1 + \sum_{l=1}^L d_l \lambda^l}. \quad (11)$$

The coefficients c_n and d_l are functions of fixed-order coefficients $f_1, f_2, \dots$. As the two coefficients f_1 and f_2 are determined, it is possible to construct two distinct Padé approximants, which reproduce Eq.(10) up to order of λ^3 :

$$\begin{aligned} R_{[1,2]} &= \frac{1 + \xi\lambda}{1 + \xi\lambda - f_2\lambda^2}, \\ R_{[2,1]} &= \frac{1 + \xi\lambda + f_2\lambda^2}{1 + \xi\lambda}, \end{aligned} \quad (12)$$

respectively. Note that, as Eq.(9) contains no term of order λ , the coefficients c_1 and d_1 are fixed to the same value:

$$\xi = -\frac{f_3}{f_2} = \frac{2(3 + \sqrt{2})}{3\pi}. \quad (13)$$




Figure 5: *Plots of fixed-order perturbative result, and Padé approximants $R_{[1,2]}$ and $R_{[2,1]}$.*

We have plotted the two Padé approximants in Fig. 5 and have made comparison with the perturbative result. Note that, thanks to the fact that f_3 is positive, both forms of the

Padé approximant show *monotonically decreasing* behavior as the ‘t Hooft coupling parameter becomes strong. This is to be contrasted with the behavior of the fixed-order perturbative result, in which the strong coupling behavior is dominated by the $f_3 g_{\text{YM}}^3$ term. Hence, within the simple Padé-improvement of the perturbation series, we have obtained a behavior consistent with the strong coupling expansion drawn from the AdS_5 supergravity analysis and hence with smooth interpolation between weak and strong coupling regimes.

Although the Padé approximant would not be able to draw any definite conclusion regarding the strong coupling behavior, at least one can say that the positive contribution of the $f_3 g_{\text{YM}}^3$ term to the free-energy density is not as disastrous as it may appear for a smooth interpolation over the ‘t Hooft coupling parameter space. Explicit calculation of higher-order g_{YM}^4 , $g_{\text{YM}}^4 \ln g_{\text{YM}}^2$ and g_{YM}^5 might be helpful in drawing more definitive conclusion on this issue.

4 Remark on D1+D5 and AdS_3 Black Hole Free-Energy

Finally, we would like to bring up an issue regarding another interesting configuration exhibiting AdS/CFT correspondence: a bound-state of N_1 D1-branes and N_5 D5-branes. Effective description of the configuration is controlled by ‘t Hooft coupling parameters, $g_{\text{YM}}^2 N_1, g_{\text{YM}}^2 N_5$.

At strong coupling limit, the bound-state of extremal D1-D5 brane is described by Type IIB string on $AdS_3 \times S^3 \times T^4$. An important observation of [3] was that the geometry is not modified even if the D1-D5 branes become *non-extremal*. This is because of the fact that the strong coupling expansion (viz. in inverse powers of the two ‘t Hooft coupling parameters) is expressed in power-series of Weyl curvatures and that the Weyl curvature vanishes for a product of AdS_3 and S^3 for equal radii of curvature. This implies that the free-energy density of the Schwarzschild AdS_3 black hole (BTZ black hole) is given, to all orders in strong coupling expansion, by

$$F = -\pi N_1 N_5 T^2, \quad (14)$$

viz. *independent* of the ‘t Hooft coupling parameters.

At weak coupling limit, the D1-D5 brane configuration is described by $d = 2$ $\mathcal{N} = 4$ supersymmetric Yang-Mills theory with gauge group $SU(N_1) \times SU(N_5)$ and bi-fundamental matters [22]. Higgs-branch of the theory is described by four Goldstone bosons living on an effectively $N_1 N_5$ times bigger spatial volume. Thus, thermodynamic free-energy may be calculated in terms of sigma model of the four Goldstone bosons. At leading order in derivative expansion, the free-energy density ought to be given by

$$F = 4 \cdot \left(-\frac{\pi}{4} T^2 \right) \cdot N_1 N_5 + \cdots, \quad (15)$$

where the ellipses denote corrections from higher-order interaction terms of the sigma model.

The leading-order term in Eq.(15) matches with the exact result of Eq.(14), so if the free-energy density were to interpolate monotonically between weak and strong coupling regimes, the free-energy density Eq.(15) should be also an exact result of perturbative expansion in powers of 't Hooft coupling parameters. This implies that, in thermodynamics of $d = 2$ super Yang-Mills theory, there exists a curious non-renormalization theorem (not related to supersymmetry). We suspect this may have to do with subtle noncommutativity between 't Hooft limit, $N_1, N_5 \rightarrow \infty$, and thermodynamic infinite volume limit, $V \rightarrow \infty$, in two dimensional gauge theories. Whether this is indeed the case would serve as an interesting checkpoint to the generality of the anticipated interpolation. Work along this direction is in progress and will be reported elsewhere.

References

- [1] J. Maldacena, Adv. Theor. Math. Phys. **2** (1998) 231.
- [2] M. Li, *Evidence for Large N Phase Transition in $\mathcal{N} = 4$ Super Yang-Mills Theory at Finite Temperature*, hep-th/9807196.
- [3] S.S. Gubser, I.R. Klebanov and A.A. Tseytlin, Nucl. Phys. **B534** (1998) 202.
- [4] S.W. Hawking and D.N. Page, Comm. Math. Phys. **87** (1983) 577.
- [5] S.-J. Rey and J.-T. Yee, *Macroscopic String as Heavy Quark in Large N Gauge Theory and Anti-de Sitter Supergravity*, hep-th/9803001.
- [6] J. Maldacena, Phys. Rev. Lett. **80** (1988) 4859, hep-th/9903002.
- [7] E. Witten, Adv. Theor. Math. Phys. **2** (1998) 505, hep-th/9803131.
- [8] S.-J. Rey, S. Theisen and J.-T. Yee, Nucl. Phys. **B527** (1998) 171, hep-th/9803135.
- [9] A. Brandhuber, N. Itzhaki, J. Sonnenschein and S. Yankielowicz, Phys. Lett. **B434** (1998) 36, hep-th/9803137.
- [10] S.S. Guber, I.R. Klebanov and A.M. Polyakov, Phys. Lett. **428B** (1998) 105, hep-th/9802109.
- [11] E. Witten, Adv. Theor. Math. Phys. **2** (1998) 253, hep-th/9802150.
- [12] S.-J. Rey, *Quarks, Mesons and Baryons out of Anti-de Sitter Superstring Theory*, to appear in the proceedings of '13th Nishinomiya Yukawa Memorial Symposium' (November, 1998).

- [13] A. Fotopoulos and T. R. Taylor, Phys. Rev. **D59** (1999) 044035, [hep-th/9811224](#).
- [14] M.A. Vázquez-Mozo, *A Note on Supersymmetric Yang-Mills Thermodynamics*, [hep-th/9905030](#).
- [15] D.J. Gross, R. Pisarski and L.G. Yaffe, Rev. Mod. Phys. **53** (1983) 43.
- [16] See, for example, J. I. Kapusta, “*Finite Temperature Field Theory*,” (Cambridge University Press, Cambridge, 1989).
- [17] P. Arnold and C. Zhai, Phys. Rev. **D50** (1994) 7603 [hep-th/9408276](#); Phys. Rev. **D51** (1995) 1906, [hep-th/9410360](#).
- [18] B. Kastening and C. Zhai Phys. Rev. **D52**, 7232 (1995), [hep-ph/9507380](#);
E. Braaten and A. Nieto, Phys. Rev. **D53**, 3421 (1996), [hep-ph/9510408](#).
- [19] See, for example, C.M. Bender and S.A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers*, (McGraw-Hill, New York, 1977), G.A. Baker and P. Graves-Morris, *Padé Approximants*, 2nd ed. (Cambridge Univ. Press, Cambridge, 1996).
- [20] S.J. Brodsky, J. Ellis, E. Gardi, M. Karliner and M.A. Samuel, Phys. Rev. **D56** (1997) 6980; A.L. Kataev and V.V. Starshenko, Mod. Phys. Lett. **A10** (1995) 235; E. Gardi, Phys. Rev. **D56** (1997) 68; I. Jack, D.R.T. Jones and M.A. Samuel, Phys. Lett. **B407** (1997) 143.
- [21] B. Kastening, Phys. Rev. **D56**, 8107 (1997), [hep-ph/9708219](#);
T. Hatsuda, Phys. Rev. **D56**, 8111 (1997), [hep-ph/9708257](#).
- [22] C.G. Callan and J. Maldacena, Nucl. Phys. **B472** (1996) 591, [hep-th/9602043](#).